

Nitrogen Rate and Timing on Corn

Study ID: 401155201702

County: Saunders

Soil Type: Aksarben silty clay loam 0-2% slope;
Judson silt loam 2-6% slopes

Planting Date: 5/9/17

Harvest Date: 11/3/17

Population: 26,500

Row Spacing (in): 30

Hybrid: NK 73Y-3111

Reps: 4

Previous Crop: Soybean

Tillage: No-Till

Herbicides: *Pre:* 2.5 qt/ac Acuron™ and 1 qt/ac

Roundup PowerMAX® on 5/10/17 *Post:* 1 qt/ac

Roundup PowerMAX®, and 1 pt/ac Dual II

Magnum® on 6/10/17

Seed Treatment: Avicta®, CruiserMaxx®

Foliar Insecticides: None

Foliar Fungicides: None

Fertilizer: 150 lb/ac of Phos Plus 36D (11-30-0-9S-

2Zn) in Dec. 2016; 2 Ton/Ac Ag Lime (90% CCE) in Jan. 2017; 4 gal/ac 12-0-0-26S, 1.5 gal/ac 9% Zn, 1.5 gal/ac 10% Boron, 0.5 gal/ac 6% Mn on 5/9/17, in addition to N applied as treatments in this study

Note: Severe wind events in late October

Irrigation: None

Rainfall (in):



Soil Sample (Oct. 2016):

OM	P1	P2	K	Mg	Ca	Na	pH	Buffer pH	C.E.C.	K	Mg	Ca	H	Na	S	Zn	Mn	Fe	Cu	B
%	-----ppm-----						--% base saturation--				-----ppm-----						-----ppm-----			
3.1	11	17	213	356	2248	20	5.8	6.6	18.3	3	16.2	61.4	18.9	0.5	15	1	10	53	0.9	0.4

Introduction: The purpose of this study was to evaluate different N management strategies on corn. Applying a portion of the N fertilizer during the growing season allows fertilizer availability to better match the time the corn is uptaking N. The N in this study was applied as 32 percent UAN. There were four treatments: 1) 140 lb/ac N as pre-plant, 2) 100 lb/ac N pre-plant plus 40 lb/ac N sidedress, 3) 100 lb/ac N pre-plant plus 40 lb/ac N sidedress and Hydras-Hume humic acid at a rate of 3 gal/ac per 100 gal of 32 percent UAN, and 4) 100 lb/ac N pre-plant plus 75 lb/ac N sidedress. The pre-plant application was on 5/9/17. The sidedress applications were made on 6/20/17 with a homemade Y-Drop type applicator. There was a 1.25" rain on June 28.

Results:

	Test Weight	Moisture (%)	Yield (bu/ac)†	Marginal Net Return‡ (\$/ac)
140 lb/ac Pre-plant N	56 A*	13.5 A	193 A	542.59 A
100 lb/ac Pre-plant N + 40 lb/ac Sidedress N	56 A	13.6 A	195 A	540.66 A
100 lb/ac Pre-plant N + 40 lb/ac Sidedress N + Humic Acid	56 A	13.6 A	199 A	549.40 A
100 lb/ac Pre-plant N + 75 lb/ac Sidedress N	56 A	13.6 A	200 A	542.61 A
P-Value	0.420	0.785	0.463	0.946

*Values with the same letter are not significantly different at a 90% confidence level.

†Bushels per acre corrected to 15.5% moisture.

‡Marginal net return based on \$3.15/bu corn, \$1.46/gal 32% liquid fertilizer (\$0.36/lb N), and \$7/ac per application.

Summary: There was no difference in test weight, moisture, yield, or net return between the treatments studied.

Sponsored by:



In Partnership with:



Nebraska Dry Bean
Commission